Table of common feed stuffs and problems

Feed	problem	solution
	1	
Paddy	Too high in fibre,	Feed only to adult
5 1 1 1 1 1	Irritates intestines	poultry and limit intake
Broken polished rice	Chicks over eat and crop	Limit intake
	become impacted.	
	Very low fibre and	
	protein	
Rice Bran and polishings	High fibre	Limit intake
	High fat, becomes rancid	Store short time
	High in bacteria	Cook for young pigs
Rice husk mixed in	Young animal irritated	Sieve low quality bran
	GIT	
Taro , bulb, leaf and stem	Oxalic acid crystals	cook
Banana stem	High fibre, low protein	Limit intake
Corn	Large hard seed	Crush for chicks and pigs
		1 6
	Medium High oil	Limit intake
	1 10 011 011	
2.5		
Most types of beans	Toxins and protein	cook
	uptake inhibitors	
Most types of beans soybeans	uptake inhibitors As above and very high	cook Cook and limit intake
soybeans	uptake inhibitors	Cook and limit intake
	uptake inhibitors As above and very high	
soybeans	uptake inhibitors As above and very high in fat	Cook and limit intake
soybeans	uptake inhibitors As above and very high in fat	Cook and limit intake Cook and soak in at least
soybeans Velvet beans	uptake inhibitors As above and very high in fat L-dopa	Cook and limit intake Cook and soak in at least two changes of water
soybeans Velvet beans	uptake inhibitors As above and very high in fat L-dopa	Cook and limit intake Cook and soak in at least two changes of water Used only in very small
soybeans Velvet beans	uptake inhibitors As above and very high in fat L-dopa toxic	Cook and limit intake Cook and soak in at least two changes of water Used only in very small percentage after
soybeans Velvet beans Jack bean	uptake inhibitors As above and very high in fat L-dopa	Cook and limit intake Cook and soak in at least two changes of water Used only in very small percentage after extensive processing Crushed and moistened.
soybeans Velvet beans Jack bean	uptake inhibitors As above and very high in fat L-dopa toxic Cyanide	Cook and limit intake Cook and soak in at least two changes of water Used only in very small percentage after extensive processing Crushed and moistened. exposed to the air for one
soybeans Velvet beans Jack bean	uptake inhibitors As above and very high in fat L-dopa toxic	Cook and limit intake Cook and soak in at least two changes of water Used only in very small percentage after extensive processing Crushed and moistened. exposed to the air for one hour
soybeans Velvet beans Jack bean	uptake inhibitors As above and very high in fat L-dopa toxic Cyanide	Cook and limit intake Cook and soak in at least two changes of water Used only in very small percentage after extensive processing Crushed and moistened. exposed to the air for one hour Limit intake mix with
soybeans Velvet beans Jack bean	uptake inhibitors As above and very high in fat L-dopa toxic Cyanide Very low protein	Cook and limit intake Cook and soak in at least two changes of water Used only in very small percentage after extensive processing Crushed and moistened. exposed to the air for one hour Limit intake mix with high protein feeds
soybeans Velvet beans Jack bean Casssava root	uptake inhibitors As above and very high in fat L-dopa toxic Cyanide Very low protein Flour dusty	Cook and limit intake Cook and soak in at least two changes of water Used only in very small percentage after extensive processing Crushed and moistened. exposed to the air for one hour Limit intake mix with high protein feeds moisten
soybeans Velvet beans Jack bean	uptake inhibitors As above and very high in fat L-dopa toxic Cyanide Very low protein	Cook and limit intake Cook and soak in at least two changes of water Used only in very small percentage after extensive processing Crushed and moistened. exposed to the air for one hour Limit intake mix with high protein feeds